



Nuclear Starbursts in Nearby Seyfert Galaxies

A. Romano^{1,2}, A. Rifatto², S. Ciroi¹, P. Rafanelli¹, M. Dopita³

¹ *Dipartimento di Astronomia, Università di Padova, vicolo dell'Osservatorio 2, 35122 Padova, Italy*

² *Osservatorio Astronomico di Capodimonte, via Moiariello 16, 80131 Napoli, Italy*

³ *Research School of Astronomy and Astrophysics, Australian National University, Private Bag, Weston Creek Post Office, ACT 2611, Australia*

ABSTRACT

Active galaxies are often associated with starbursts, which are possible sources of fuelling for the nuclear black holes. By means of optical (3500 – 7500 Å) spectroscopic and narrow-band photometric data ($H\alpha$ + $[NII]\lambda 6548, 6583$ and $[OIII]\lambda 5007$) it is possible to detect starburst features in active nuclei and to study the correlation with Seyfert type and the surrounding environment. Here we present evidences of young stellar populations (HII high order Balmer lines) and extended $H\alpha$ emission (indicative of HII regions) in 6 nearby Seyfert galaxies ($z < 0.02$). These very preliminary results seem to suggest that the nuclear starburst presence does not depend on the Seyfert type, favouring the hypothesis of the AGN Unified Model. By estimating the present SFR ($< 10^6$ years) and recent SFR ($< 10^9$ years) we observe that these objects have experienced most of star formation episodes in past epochs. This supports the hypothesis of a starburst-AGN connection, favouring an evolutionary scenario in which the AGN survives to the starburst.

1. INTRODUCTION

One of the most important questions concerning the Active Galactic Nuclei (AGNs) phenomenon is the connection between starburst and nuclear activity. The role of starbursts in AGNs has been extensively discussed in the past and several theoretical scenarios have been proposed, from those in which the two phenomena are indirectly connected because both are triggered by and live on gas fuelling (e.g., mergers, interactions between galaxies, bars, etc.) to those in which the starbursts are directly bound with the AGNs as possible sources of fuelling for the nuclear black holes. The study of the extended circum- and extra-nuclear regions of ionized gas in active galaxies can shed more light on the interaction between the active nucleus and its environment, because the gas is a convenient tracer of massive star-formation (HII regions), ionization by non thermal processes (e.g., power-law continua), and/or shocks. Evidences of starburst features have been detected in numerous Seyfert 2 (e.g. Cid Fernandes et al. 2001), but rarely in Seyfert 1 galaxies (Rodríguez-Ardila et al. 2003). On the contrary, if the AGNs Unified Model (Antonucci,




Fig. 1. Upper panel: The $H\alpha$ (left) and $[OIII]\lambda 5007$ (right) emission-line images of ESO018-G09. Bottom panel: Section of the nuclear spectrum of ESO018-G09 (Seyfert 2), where we see the absorption lines of young stars.

1993) is valid and a connection exists between AGN and starburst, one would expect to observe a similar distribution of circumnuclear star-forming regions among the different types of AGNs. In this poster we present preliminary results of an investigation of the nuclear and circumnuclear regions of 6 nearby Seyfert galaxies. Narrow-band images isolating the emission lines of $H\alpha + [NII]\lambda 6548, 6583$ and $[OIII]\lambda 5007$ were used to search for spatially extended circumnuclear emission regions. These observations were followed up with long-slit, medium resolution spectrophotometry to probe further the ionization state of the gas. The studied galaxies involve all Seyfert types (three Seyfert 2, one Seyfert 1, one Seyfert 1.5 and one Seyfert 1.8) in order to check a possible correlation between the presence of starburst and the Seyfert type.




Fig. 2. Upper panel: The $H\alpha$ (left) and $[OIII]\lambda 5007$ (right) emission-line images of ESO362-G18 (Seyfert 1.8). Bottom panel: The nuclear spectrum of ESO362-G18, where the absorption lines of young stars are detectable.

2. THE OBSERVATIONS

The photometric observations were carried out in October 2001 and January 2002 at the Siding Spring Observatory (Australia) with 2.3m telescope. A 1kx1k CCD camera (pixel size = $24 \mu m$, scale = $0.6''/px$) was used in combination with $H\alpha + [NII]\lambda 6548, 6583$ and $[OIII]\lambda 5007$ filters (FWHM $\sim 75 \text{ \AA}$). Long-slit spectra of 4 out of 6 galaxies (ESO018-G09, ESO362-G18, ESO377-G24, ESO428-G14) were obtained at the same telescope with the Dual Beam Spectrograph (pixel size = $15 \mu m$, scale = $0.89''/px$), which allowed to obtain simultaneously the blue (3640-5550 $\text{\AA}$) and the red (5540-7450 $\text{\AA}$) parts of the optical wavelength range at medium resolution ($\sim 2 \text{ \AA}$). The spectra of the other 2 galaxies were obtained in December 1990 and March 1991 at ESO-La Silla (Chile) at the 1.52m telescope with the B&C Spectrograph (pixel size = $15 \mu m$, scale = $0.82''/px$) at lower resolution. The covered spectral range was 4620-7470 $\text{\AA}$ for NGC 3081 and 3730-6780 $\text{\AA}$ for NGC 1365. The data were reduced and analyzed with the IRAF software.

All the observations were bias subtracted, flat field corrected, cleaned from cosmic rays and sky subtracted. The emission-line images were calibrated in flux using a planetary nebula observed in the same runs with the same instrumental configuration. The two-dimensional spectra were calibrated in wavelength using a Ne-Ar lamp for SSO spectra and He-Ar lamp for ESO spectra, then calibrated in flux using a spectrophotometric standard star. For each galaxy we extracted spectra of the nucleus and several circum- and/or extra-nuclear regions, whose sizes were determined by means of both the $H\alpha$ line profile along the slit and the extension of the emission in the $H\alpha$ images. We identified the stellar absorption lines and bands in each spectrum. Then, the template of a S0 galaxy was subtracted to remove the underlying stellar population contribution. After this step, we identified the emission lines in the spectra of each object: their profiles were decomposed into multiple Gaussians using the IRAF task FITPROF. In particular, we measured the redshift, width and flux of each identified line. Then, the line fluxes were corrected for internal reddening using the observed Balmer decrement $H\alpha/H\beta$ chosen equal to 3.1 for the active nuclei and 2.85 for the starburst regions (Osterbrock, 1989), and applying the extinction curve by Cardelli, Clayton & Mathis (1989).

3. RESULTS

We found clear evidences of the presence of young stellar populations (HI high order Balmer absorption lines (HOBs), indicative of age < 1 Gyr) and extended $H\alpha$ emission (indicative of HII regions) in 4 galaxies of the sample: a Seyfert 2, a Seyfert 1, a Seyfert 1.5 and a Seyfert 1.8 (see figures 1-2). These preliminary results seem to suggest that the nuclear starburst presence does not depend on the Seyfert type, favouring the hypothesis of the AGN Unified Model. We also estimated the present SFR (on time-scale of 10^6 years) and compared it with the recent SFR (on time-scale of 10^9 years) for every object of the sample. We observed that the galaxies which do not show clear signs of present starbursts have had major star formation in the past epochs. This result supports the hypothesis of a starburst-AGN connection, favouring an evolutionary scenario in which the AGN survives to the starburst. In this scenario we should expect that the composed Seyfert+starburst system evolves into a pure Seyfert. Moreover, we observed that the presence of nuclear starburst features is usually associated to the presence of extended star formation regions spread over the disk. Given the small number of the studied objects, we cannot suggest any hypothesis at this stage about the role of the interactions in the starburst-AGN connection. But our future goals are just to extend this analysis to a larger sample of isolated Seyfert galaxies in order to verify these preliminary results on a statistically significant sample.

Acknowledgements. This research was partially based on data from a INAF-OAC and ANU-MSSSO project, in collaboration with the Department of Astronomy, University of Padova.

REFERENCES

- Antonucci, R. 1993, ARA&A, 31, 473
 Cardelli, J. A., Clayton, G. C. & Mathis, J. S. 1989, ApJ, 345, 245
 Cid Fernandes, R., Heckman, T., Schmitt, H., Delgado, R.M.G. & Storchi-Bergmann, T. 2001, ApJ, 558, 81
 Osterbrock, D. E. 1989, Astrophysics of Gaseous Nebulae and AGN, Oxford University Press
 Rodriguez-Ardila, A. & Viegas, S. M. 2003, MNRAS, 340, L33